

Contents

| | |
|---|-----|
| Preface | 1 |
| In Memoriam | 3 |
| Abstracts | 5 |
| Genetic markers and their application | |
| The development and application of genetic markers for the Kuruma prawn <i>Penaeus japonicus</i> S.S. Moore, V. Whan, G.P. Davis, K. Byrne, D.J.S. Hetzel and N. Preston (Queensland, Australia) | 19 |
| Linkage analysis of quantitative trait loci associated with spawning time in rainbow trout (<i>Oncorhynchus mykiss</i>) T. Sakamoto (Guelph, Ont., Canada; Tokyo, Japan), R.G. Danzmann (Guelph, Ont., Canada), N. Okamoto (Tokyo, Japan), M.M. Ferguson (Guelph, Ont., Canada) and P.E. Ihssen (Maple, Ont., Canada) | 33 |
| Epistasis in allelic expression at upper temperature tolerance QTL in rainbow trout R.G. Danzmann, T.R. Jackson and M. M. Ferguson (Guelph, Ont., Canada) | 45 |
| Cosmid clones from Atlantic salmon: physical genome mapping M. Lundin, B. Mikkelsen (Oslo, Norway), P. Moran, J.L. Martinez (Oviedo, Spain) and M. Syed (Oslo, Norway) | 59 |
| Identification of a sex-linked GH pseudogene in one of two species of Japanese salmon (<i>Oncorhynchus masou</i> and <i>O. rhodurus</i>) I. Nakayama (Tamaki, Japan), C.A. Biagi (Vancouver, Canada), N. Koide (Vancouver, Canada; Hokkaido, Japan) and R.H. Devlin (Vancouver, Canada) | 65 |
| Restriction fragment length polymorphism at the growth hormone 1 gene in Atlantic salmon (<i>Salmo salar</i> L.) and its association with weight among the offspring of a hatchery stock R. Gross (Tartu, Estonia) and J. Nilsson (Umeå, Sweden) | 73 |
| Identification of candidate DNA markers associated with IHN virus resistance in backcrosses of rainbow (<i>Oncorhynchus mykiss</i>) and cutthroat trout (<i>O. clarki</i>) Y. Palti (Pullman, WA, USA), J.E. Parsons (Twin Falls, ID, USA) and G.H. Thorgaard (Pullman, WA, USA) | 81 |
| Genetic identification of sparid species by isozyme markers: application to interspecific hybrids J.A. Alarcón and M.C. Alvarez (Málaga, Spain) | 95 |
| Breeding and selective improvement | |
| Early growth performance of Atlantic salmon full-sib families reared in single family tanks versus in mixed family tanks C.M. Herlinger, P.T. O'Reilly, R.W. Doyle, J.M. Wright (Halifax, Canada) and F. O'Flynn (New Brunswick, Canada) | 105 |
| Outbreeding depression in hybrids between odd- and even-broodyear pink salmon A.J. Gharrett, W.W. Smoker (Juneau, AK, USA), R.R. Reisenbichler (Seattle, WA, USA) and S.G. Taylor (Juneau, AK, USA) | 117 |

Contents

| | |
|---|-----|
| Preface | 1 |
| In Memoriam | 3 |
| Abstracts | 5 |
| Genetic markers and their application | |
| The development and application of genetic markers for the Kuruma prawn <i>Penaeus japonicus</i> S.S. Moore, V. Whan, G.P. Davis, K. Byrne, D.J.S. Hetzel and N. Preston (Queensland, Australia) | 19 |
| Linkage analysis of quantitative trait loci associated with spawning time in rainbow trout (<i>Oncorhynchus mykiss</i>) T. Sakamoto (Guelph, Ont., Canada; Tokyo, Japan), R.G. Danzmann (Guelph, Ont., Canada), N. Okamoto (Tokyo, Japan), M.M. Ferguson (Guelph, Ont., Canada) and P.E. Ihssen (Maple, Ont., Canada) | 33 |
| Epistasis in allelic expression at upper temperature tolerance QTL in rainbow trout R.G. Danzmann, T.R. Jackson and M. M. Ferguson (Guelph, Ont., Canada) | 45 |
| Cosmid clones from Atlantic salmon: physical genome mapping M. Lundin, B. Mikkelsen (Oslo, Norway), P. Moran, J.L. Martinez (Oviedo, Spain) and M. Syed (Oslo, Norway) | 59 |
| Identification of a sex-linked GH pseudogene in one of two species of Japanese salmon (<i>Oncorhynchus masou</i> and <i>O. rhodurus</i>) I. Nakayama (Tamaki, Japan), C.A. Biagi (Vancouver, Canada), N. Koide (Vancouver, Canada; Hokkaido, Japan) and R.H. Devlin (Vancouver, Canada) | 65 |
| Restriction fragment length polymorphism at the growth hormone 1 gene in Atlantic salmon (<i>Salmo salar</i> L.) and its association with weight among the offspring of a hatchery stock R. Gross (Tartu, Estonia) and J. Nilsson (Umeå, Sweden) | 73 |
| Identification of candidate DNA markers associated with IHNV resistance in backcrosses of rainbow (<i>Oncorhynchus mykiss</i>) and cutthroat trout (<i>O. clarki</i>) Y. Palti (Pullman, WA, USA), J.E. Parsons (Twin Falls, ID, USA) and G.H. Thorgaard (Pullman, WA, USA) | 81 |
| Genetic identification of sparid species by isozyme markers: application to interspecific hybrids J.A. Alarcón and M.C. Alvarez (Málaga, Spain) | 95 |
| Breeding and selective improvement | |
| Early growth performance of Atlantic salmon full-sib families reared in single family tanks versus in mixed family tanks C.M. Herlinger, P.T. O'Reilly, R.W. Doyle, J.M. Wright (Halifax, Canada) and F. O'Flynn (New Brunswick, Canada) | 105 |
| Outbreeding depression in hybrids between odd- and even-broodyear pink salmon A.J. Gharrett, W.W. Smoker (Juneau, AK, USA), R.R. Reisenbichler (Seattle, WA, USA) and S.G. Taylor (Juneau, AK, USA) | 117 |

| | |
|---|-----|
| Variation in development rate among clonal lines of rainbow trout (<i>Oncorhynchus mykiss</i>) B.D. Robison, P.A. Wheeler and G.H. Thorgaard (Pullman, WA, USA) | 131 |
| Responses to two generations of index selection in Atlantic salmon (<i>Salmo salar</i>) F.M. O'Flynn (Kerry, Ireland; New Brunswick, Canada), J.K. Bailey and G.W. Friars (New Brunswick, Canada) | 143 |
| Genotype-environment interactions in the susceptibility of the common carp, <i>Cyprinus carpio</i> , to <i>Ichthyophthirius multifiliis</i> infections D.J. Price and G.M. Clayton (Plymouth, UK) | 149 |
| Growth of five Asian red tilapia strains in saline environments M.R.R. Romana-Eguia and R.V. Eguia (Binangonan, Philippines) | 161 |
| Strain evaluation of striped bass (<i>Morone saxatilis</i>) under controlled conditions J.M. Jacobs (Cambridge, MD, USA), S. Lindell (Turners Falls, MA, USA), W. Van Heukelem (Cambridge, MD, USA), E.M. Hallerman (Blacksburg, VA, USA) and R.M. Harrell (Cambridge, MD, USA) | 171 |
| Definition of a breeding objective for commercial production of the freshwater crayfish, marron (<i>Cherax tenuimanus</i>) M. Henryon (Tjele, Denmark), I.W. Purvis (Armidale, Australia) and P. Berg (Tjele, Denmark) | 179 |
| Stocking density and tank size in the design of breed improvement programs for body size of tilapia G.A.E. Gall and Y. Bakar (California Davis, CA, USA) | 197 |
| Sex determination and its manipulation | |
| Evidence for the presence of sex chromosomes in the Nile tilapia (<i>Oreochromis niloticus</i>) from synaptonemal complex analysis of XX, XY and YY genotypes L.A.P. Carrasco, D.J. Penman and N. Bromage (Stirling, UK) | 207 |
| Environmental sex determination: the effect of temperature and salinity on sex ratio in <i>Oreochromis niloticus</i> L. J.S. Abucay (Nueva Ecija, Philippines), G.C. Mair, D.O.F. Skibinski and J.A. Beardmore (Swansea, UK) | 219 |
| Studies on sex-manipulation and production of cloned populations in hirame, <i>Paralichthys olivaceus</i> (Temminck et Schlegel) E. Yamamoto (Tottori, Japan) | 235 |
| Monosex female production in the silver barb (<i>Puntius gonionotus</i> Bleeker) N. Pongthana (Pathum' Thani, Thailand), D.J. Penman (Stirling, UK), P. Baoprasertkul (Pathum' Thani, Thailand), M.G. Hussain, M. Shahidul Islam (Mymensingh, Bangladesh), S.F. Powell and B.J. McAndrew (Stirling, UK) | 247 |
| Sex determination and the feasibility of genetically male tilapia production in the Thai-Chitralada strain of <i>Oreochromis niloticus</i> (L.) P.A. Tuan (Bac Ninh, Viet Nam; Swansea, UK), G.C. Mair (Swansea, UK), D.C. Little (Klong Luang, Thailand) and J.A. Beardmore (Swansea, UK) | 257 |
| Transgenesis | |
| Mendelian transmission, transgene dosage and growth phenotype in transgenic tilapia (<i>Oreochromis hornorum</i>) showing ectopic expression of homologous growth hormone R. Martínez, A. Arenal, M.P. Estrada, F. Herrera, V. Huerta, J. Vázquez, T. Sánchez and J. De la Fuente (Havana, Cuba) | 271 |
| Growth performance studies in transgenic <i>Cyprinus carpio</i> Y. Hinitz and B. Moav (Ramat Aviv, Israel) | 285 |
| Effect of an osmotic differential on the efficiency of gene transfer by electroporation of fish spermatozoa J.-H. Kang, G. Yoshizaki, O. Homma, C.A. Strüssmann and F. Takashima (Tokyo, Japan) | 297 |

| | |
|--|-----|
| A decision support software for safely conducting research with genetically modified fish and shellfish E. Hallerman, D. King (Blacksburg, VA, USA) and A. Kapuscinski (St. Paul, MN, USA) . . . | 309 |
| Transfer and expression of glucose transporter and hexokinase genes in salmonid fish T.I. Pitkänen, A. Krasnov, M. Reinisalo and H. Mölsä (Kuopio, Finland) | 319 |
| Growth performance of transgenic tilapia containing an exogenous piscine growth hormone gene M.A. Rahman and N. Maclean (Southampton, UK) | 333 |
| Chromosome set manipulation | |
| Aneuploid Pacific oyster (<i>Crassostrea gigas</i> Thunberg) as incidentals from triploid production Z. Wang (Shandong, China), X. Guo, S.K. Allen Jr. (Port Norris, NJ, USA) and R. Wang (Shandong, China) | 347 |
| Genetic variation in growth parameters until commercial size in diploid and triploid freshwater rainbow trout (<i>Oncorhynchus mykiss</i>) and seawater brown trout (<i>Salmo trutta</i>) S. Bonnet, P. Haffray (Rennes, France), J.M. Blanc, F. Vallée (Saint-Pée-sur-Nivelle, France), C. Vauchez (Rennes, France), A. Fauré (Sizun, France) and B. Fauconneau (Rennes, France) | 359 |
| Proof of the successful development of Nile tilapia (<i>Oreochromis niloticus</i>) clones by DNA fingerprinting I. Jenneckens, A. Müller-Belecke, G. Hörstgen-Schwark and J.-N. Meyer (Göttingen, Germany) | 377 |
| Growth and gonadal development in triploid sea bass (<i>Dicentrarchus labrax</i> L.) during the first two years of age A. Felip, S. Zanuy, M. Carrillo (Castellón, Spain) and F. Piferrer (Barcelona, Spain) | 389 |
| Induction of diploid gynogenesis using interspecific sperm and production of tetraploids in African catfish, <i>Clarias gariepinus</i> Burchell (1822) L. Várad, I. Benkó, J. Varga and L. Horváth (Páter, Hungary) | 401 |
| Genetic resources, conservation and management | |
| Genetic variability and pedigree tracing of a hatchery-reared stock of red sea bream (<i>Pagrus major</i>) used for stock enhancement, based on microsatellite DNA markers R. Perez-Enriquez, M. Takagi and N. Taniguchi (Nankoku City, Japan) | 413 |
| Genetic differences among year classes in a hatchery population of coho salmon (<i>Oncorhynchus kisutch</i> (Walbaum, 1792)) in Chile F.M. Winkler (Coquimbo, Chile), D. Bartley (Rome, Italy) and N.F. Díaz (Santiago, Chile) . . | 425 |
| Genetic variability of German and foreign common carp (<i>Cyprinus carpio</i> L.) populations K. Kohlmann and P. Kersten (Berlin, Germany) | 435 |
| Population genetic structure of lobster (<i>Homarus gammarus</i>) in Norway, and implications for enhancement and sea-ranching operation K.E. Jørstad and E. Færevit (Bergen, Norway) | 447 |
| Genetic differentiation between Tasmanian cultured Atlantic salmon (<i>Salmo salar</i> L.) and their ancestral Canadian population: comparison of microsatellite DNA and allozyme and mitochondrial DNA variation A. Reilly, N.G. Elliott, P.M. Grewe (Hobart, Australia), C. Clabby (Hull, UK), R. Powell (University College Galway, Ireland) and R.D. Ward (Hobart, Australia) | 459 |
| Genetic resources of commercially important fish species in the Czech Republic: present state and future strategy M. Flajšhans, O. Linhart (Vodňany, Czech Republic), V. Šlechtová and V. Šlechta (Liběchov, Czech Republic) | 471 |
| Genetic diversity among wild forms and cultivated varieties of <i>Discus</i> (<i>Symphysodon</i> spp.) as revealed by Random Amplified Polymorphic DNA (RAPD) fingerprinting T.L. Koh, G. Khoo, L.Q. Fan and V.P.E. Phang (Kent Ridge, Singapore) | 485 |

